

THE UNDERLYING LOGIC IS MANDATORY ALSO IN DISCUSSING THE PHILOSOPHY OF QUANTUM PHYSICS

DÉCIO KRAUSE

Graduate Course in Logic and Metaphysics, Federal University of Rio de Janeiro/CNPq, BRAZIL

deciokrause@gmail.com

<https://orcid.org/0000-0003-0463-2796>

Abstract. Any scientific theory (here we consider physical theories only) has an underlying logic, even if it is not totally made explicit. The role of the underlying logic of a theory T is mainly to guide the proofs and the accepted consequences of the theory's principles, mainly when described by its axioms. In this sense, the theorems of the underlying logic are also theorems of the theory. In most cases, if pressed, the scientist will say that the underlying logic of most physical theories is classical logic or some fragment of a set theory suitable for accommodating the theory's mathematical and logical concepts. We argue that no physical theory and no philosophical discussion based on the theory should dismiss its underlying logic, so the arguments advanced by some philosophers of physics in that certain entities (the considered case deals with quantum entities) can be only weakly discerned or be just 'relational' and that they cannot be *absolutely* identified by a monadic property, are fallacious once one remains within a 'standard' logico-mathematical framework, grounded on classical logic. We also discuss the (to us) unjustifiable claim that some properties (such as those that came from logic) would be 'illegitimate' for discerning these entities. Thus, this paper may interest logicians, physicists, and philosophers.

Keywords: underlying logic • weak discernibility • absolute discernibility • distinguishing quantum entities

RECEIVED: 20/11/2023

ACCEPTED: 09/08/2024

No logic can be imposed a priori to our systematisations.

Newton da Costa

1. Introduction

With some exceptions, mainly in the fields of mathematics and logic, most scientific theories are first put in an informal way, that is, they don't emerge either axiomatically or as formal theories. Thus, for instance, we may think of Galileo's theory of falling bodies, Darwin's theory of natural selection or Fourier's theory of heat transfer as posed informally. They are the product of several things, but mainly of scientists'



© 2024 The author(s). Open access under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License.

abilities and skills. In the process of the formation of such informal theories, ‘anything goes’, as Feyerabend would say.¹ But once the theory acquires a certain maturity, usually it passes by the scrutiny of rigour and foundational issues. This point of view of course does not exclude that a theory in its ‘final’ axiomatic form can arise already put this way, that is, axiomatically.

But when one is interested in the meta-study, a theory can be seen as an ordered pair $\mathcal{T} = \langle F, M \rangle$, where F is a ‘mathematical formalism’ (a formal system) and M is a class of set-theoretical structures, the *models* of the theory (Dalla Chiara and Toraldo di Francia 1981, p.60).

Of course, this is an idealisation. The underlying logic is encoded in the ‘ F ’ part of the schema.² For a discussion, at least in principle we can suppose that any theory can be written in the above way. The discussion whether the ‘ F ’ part should be written in first or in higher-order languages will be not touched here, despite its importance.³

This entails that the F theorems are theorems of \mathcal{T} , and no one should dispute this. So, if F is first-order logic encompassing an amount of set theory (say, from the ZFC system), which apparently is enough for standard physical theories, then for any theory the following are examples of theorems, being α , β formulas, and x and y individual variables (ranging over the domain of discourse):

1. $\neg\alpha, \alpha \vdash \beta$ the Explosion Rule
2. $\forall x \forall y ((\alpha(x) \leftrightarrow \alpha(y)) \rightarrow x = y)$ the Identity of Indiscernibles
3. (With the Axiom of Choice) Every set is equipotent to an ordinal.
4. In a standard set theory such as the ZFA (or ZFC) system, given any object a it can be discerned from *any other* object by the predicate we call ‘the identity of a ’, namely, $I_a(x) := x \in \{a\}$.

The Explosion Rule says that if one derives (or assumes) two contradictory formulas or sentences α and $\neg\alpha$, then *anything* β , written in the language of the theory, can be derived. Since this is a theorem of first-order logic, it is also a theorem of any theory grounded in classical logic. Even in the informal version of the theory, it is acknowledged that the presence of two contradictory sentences is bad for the theory, so we can accept that the Explosion Rule is valid even in this case.⁴

The Identity of Indiscernibles, a version of Leibniz’s Principle of the Identity of Indiscernibles (PII) (Rodriguez-Pereyra 2014) says that if anything (expressed by a formula) holding for x holds also for y , then x and y cannot denote distinct objects: they are (numerically) *identical*. By the way, this is the way identity is interpreted in standard logic: *sameness*.

The third fact entails that to any set we can associate an ordinal, and since ordinals are well-ordered sets, the elements of the set need to be distinct from each other.

Cardinals are particular ordinals, more precisely, they are those ordinals that are not equipotent to any ordinal less than themselves. For instance, the set O of odd natural numbers is equipotent to the set $\omega = \{0, 1, 2, \dots\}$ of the natural numbers (just take the bijection $f(x) = 2x + 1$ to show that). But it is also equipotent to the ordinal $\omega + 1 = \{0, 1, 2, \dots, \omega\}$; define $f : \omega + 1 \rightarrow O$ by $f(\omega) = 0$ and $f(x) = 2x + 1$ for any $x < \omega$. But since $\omega < \omega + 1$ in the ordinal ordering (membership)⁵ and the set of odd natural numbers is not equipotent to any ordinal less than ω , then ω is its cardinal ($\omega + 1$ is not a cardinal), which in the language of cardinals we call ' \aleph_0 '. This item has importance in the foundations of quantum physics; if we agree with most physicists that particles of the same kind are indiscernible but that a collection of them may have a cardinal (expressing how many systems there are), then how can we attribute a cardinal to such a collection without committing it to an ordinal, which would discern its elements? This point is not touched here despite its importance; see Krause (2023), Krause & Wajch (2023).

The fourth case is also relevant to the philosophy of quantum mechanics. It is well known that quantum theory has brought a lot of questions concerning several philosophical and foundational aspects, and this was also with logic. It is my opinion that the *logic of quantum theories*, and not only the field of quantum logics (Dalla Chiara, Giuntini & Greechie 2004) needs to be further considered. For instance, the fourth condition above says that an object a of the domain of our theory grounded in a classical setting is always *distinguishable* from any 'other' object by its identity, given by a *monadic property* defined as $I_a := x \in \{a\}$; in other words, an object is 'absolutely indistinguishable' just from itself. We shall be back to this point below. Thus, if the identity of the elements of the domain is counted among the available properties, then the identity of indiscernibles is a theorem of our theory and every object can be discerned from any other by a monadic property. In other words and just to emphasize, if we have a set $A = \{a, b\}$ whose cardinal is two, then its elements are *different* and can be discerned *absolutely*, being them representing quanta or not.

Summing up, we need to acknowledge that all logical and mathematical theorems of the theory's underlying logic are theorems of the theory also. This fact poses a challenge to some philosophers who do not pay due attention to the underlying logical facts in the considered physical theories.

From the few cases exemplified above, it is clear that the role played by the underlying logic of physical theories, with a special emphasis on quantum theory, is extremely relevant, even if such logic is taken only informally. Here we focus on the discussion of the very nature of the basic quantum entities and mainly on the third and fourth points above. The paper is organised as follows. In the next section, we show that if one does not pay attention to the underlying logic, then some mistakes may appear; the case of *relational*s in quantum theory is discussed, and in particular the case of *weakly discernibility*. In the sequence, we turn to another usually

assumed fact which is also problematic under the consideration of the underlying logic, namely, the possibility of discerning kinds of properties.

2. Careless with the underlying logic

Using ‘standard’ mathematical frameworks in the underlying logic, say a system of set theory such as ZFC or ZFA,⁶ which we can assume is grounded on first-order logic (Jech 2003), we need to acknowledge that we should not be careless with some results entailed by such systems. One of the main ones which interest us here is that *everything becomes an individual*. By an ‘individual’ we understand (according to our supposed informal account of the notion) a *something* that (i) is an *one* of a kind, say a person, a chair, a pen; (ii) it presents *identity* in the sense that it can be recognised *as such* in a certain situation and (iii) can be re-identified *as such* individual in other contexts or at different times. From a strictly logical point of view, this last condition should be qualified. Standard logic does not involve time, so we need to paraphrase the condition speaking in different contexts instead of different instants of time. If we agree with a definition of the number two, it is expected that we use this notion as the same one in different contexts. Anyway, the definition is informal, so surely the attentive reader will notice the way of speaking.

This last condition is essential; Julius Caesar was (*supposed to be*) the same person in Rome and passing the Rubicon. In other words, diachronic identity applies to individuals.⁷ This theory of identity implies that once we have *two* things, they are *different* and then some property does exist which ‘makes the difference’. There is no escape from this conclusion once we get rid of all notions related to substratum, haecceities and the like. Our ‘classical’ logic, standard mathematics and classical physics were elaborated with such kinds of entities (individuals) in our minds. A physical object, a triangle (in analytic geometry) or whatever thing we describe using such theories, are individuals in this sense.

But quantum physics (both orthodox quantum mechanics, henceforth ‘QM’ and quantum field theories) challenge this notion since quantum entities can be *completely indiscernible*, and indiscernibility, or indistinguishability, is a core concept in these disciplines; as Anthony Zee says, “[i]ndistinguishability has astounding consequences in the quantum domain” (Zee 2023, p. 300). The issue of *indistinguishability*, or *indiscernibility* (we use both words interchangeably) is a fundamental notion in this discipline and we relegate it on a pair with nonlocality, contextuality and entanglement (see de Barros *et al.* 2023). Indistinguishable things cannot be discerned, and this is (of course) the meaning of the word. But we can distinguish among several types of indistinguishabilities, and here we consider only two: *relative* indistinguishability when two or more things share *some* properties or relations,⁸ and *absolute* in-

distinguishability, when they share *all* their properties and relations, the meaning of ‘all’ being postponed by now. The other way around is also useful to be mentioned: things are *relatively discerned* when there is at least one relation which distinguishes them, and they are *relational*s when can be discerned by relations only, but not by (monadic) properties (Muller 2015). Furthermore, things are *absolutely discernible* when they can be distinguished by a monadic property.

The sample case is in quantum physics when we try to understand quantum objects.⁹ One of the most propagated claims is that which asserts that quantum objects cannot be distinguished *absolutely*, but only by relations; to some philosophers, in certain situations, such as the two electrons in a neutral Helium atom, we should acknowledge that they are only *weakly discernible*, that is, share an irreflexive but symmetric relation such as ‘to have spin opposite to’ but no monadic property can discern them (Muller & Saunders 2008). We ought to say that these assumptions ignore the underlying logic and that yes, things represented in the above mentioned ‘classical’ settings *can* be discerned absolutely, but this will depend on the setting and of our understanding of the term ‘property’. I shall consider in a moment the argument that these discerning properties given by the underlying logic would not be considered as ‘legitimate’.

The problem, as I see it, is that in general philosophers don’t make reference to the mathematical framework they are presupposing, and we should agree that they are presupposing some. Fred Muller, for instance, claims that “the ultimate constituents of physical reality are relational; that is, entities that are discernible by relations but not by properties. I defend this position.” (Muller 2015). But in his paper with Saunders (2008), where they say that fermions such as the electrons of a neutral Helium atom are just weakly discernible but not absolutely discernible, they explicitly mention that they are working within the ZFC system. Good, but not enough because in being so, they should not ignore the consequences of assuming ZFC (ZFA or any ‘standard’ mathematical framework you chose which encompasses the standard theory of identity), namely, that such a theory entails that any object it describes can be discerned ‘absolutely’ from any other.

Much (if not all) of their argument against absolute discernibility is grounded on their hypothesis that the discerning properties should have a ‘physical content’, although the ‘physical content’ cannot be achieved from purely syntactical means, that is, by logic, requiring semantics; we shall be back to this point soon. Let us analyse T. Bigaj’s discussion given at chapter 4 of his book (Bigaj 2022), adapting and simplifying the notation, since he resumes much of what is discussed in (Muller 2015, Muller & Saunders 2008) and in other papers by these authors.

Let L be a first-order language and \mathfrak{A} an interpretation with domain D . Then

1. Objects $a, b \in D$ are *absolutely discernible* ($\text{Abs}(a, b)$) iff there is an open for-

mula $\phi(x)$ in one free variable such that $\mathfrak{A} \models \phi(a)$ but $\mathfrak{A} \not\models \phi(b)$ (ibid., p.73).

2. Objects $a, b \in D$ are *relatively discernible* ($Rel(a, b)$) iff there is an open formula $\phi(x, y)$ in two free variables such that $\mathfrak{A} \models \phi(a, b)$ but $\mathfrak{A} \not\models \phi(b, a)$ (ibid., p.76). Notice that the hypothesis requires that a and b are discernible, and no criterion is given for that.
3. Objects $a, b \in D$ are *weakly discernible* ($Weak(a, b)$) iff there is a formula $\phi(x, y)$ in two free variables such that $\models \phi(a, b)$, but $\mathfrak{A} \not\models \phi(b, a)$, $\mathfrak{A} \not\models \phi(a, a)$ and $\mathfrak{A} \not\models \phi(b, b)$ (ibid., p.76). Again, this definition presupposes that a and b can be discerned and nothing is said about that.

Then Bigaj proves (2022, p.81) that

$$(1) \quad Abs(a, b) \rightarrow Rel(a, b) \rightarrow Weak(a, b),$$

but says that the converse arrows do not hold. To state that, he recurs to graphs. He simply says that “there are cases of objects that are weakly but not relatively discerned (and therefore absolutely discernible), and cases of relatively but not absolutely discernible. Examples of these cases are usually depicted in the form of graphs, where vertices (nodes) represent objects, and connecting arrows represent binary relations.”

The problem with this example is that most books on graph theory treat them as sets. By the way, I didn’t find a book that didn’t (see also Ore (1962), Wilson (1996)). In considering this, the above reasoning seems misleading. Let us consider the approach given by Bondy & Murty (2008). There, the authors (adapting the notation), say that a graph is an ordered triple (hence, a set) of the form $\mathcal{G} = \langle V, E, \psi \rangle$, where V is the *set* of vertices, E is the *set* of edges, and ψ is the incidence function (hence also a set) which associates an unordered pair $\{u, v\}$ (whose elements are not necessarily distinct) to each edge $e \in E$. The pair $\{u, v\}$ is written ‘ uv ’ and we say that the edge e *joins* the vertices u and v .

An isomorphism between two graphs $\mathcal{G} = \langle V, E, \psi \rangle$ and $\mathcal{G}' = \langle V', E', \psi' \rangle$ is an ordered pair (θ, ϕ) of functions such that $\theta : V \rightarrow V'$ and $\phi : E \rightarrow E'$, and $\psi(e) = uv$ iff $\psi'(\phi(e)) = \theta(u)\theta(v)$. It is easy to prove that an isomorphism is an equivalence relation. Then, any representative of an equivalence class of isomorphic graphs is called an *unlabelled* graph (op.cit., p.14).

As we see, even unlabelled graphs are *sets* and the elements of sets in standard set theories are absolutely discerned from one another by their identity properties. Unlabelled graphs, then, ‘ignore’ the nature of the vertices and edges, keeping with the ‘format’ of the graph only (see Figure 1). But to ignore the identity of vertices and edges does not correspond to say that these identities do not exist: really, they *do exist* in the underlying mathematics.

You could suggest that unlabelled graphs are just vertices linked by edges without any commitment to sets, as Ladyman and Ross seem to do (Ladyman & Ross 2007,




Figure 1: In (a), a labelled oriented graph and an unlabelled non-oriented graph in (b). This would be the case of relational but not absolutely discernible things. In (c), the case of weakly discernible but not relational discernible things Bigaj (2022, p.81).

§3.1), and so they could be used to model weak discernibility, as they suggest. We are not questioning that, but one should not take graphs *from nowhere* and use them in a foundational study. Really, to say that a graph is a set of vertices (perhaps) linked by some edges is to say little. Furthermore, in order to prove that the distinctions among edges and among vertices do exist, it is enough to find an embedding of the structure of graphs within an Euclidean space.

Perhaps you could find an alternative to seeing graphs as sets by resorting to other mathematical frameworks. Let me mention one at least, *multiset theory*. A multiset is a collection of things but some of them may appear more than once and be counted more than once. Thus, the multiset $A = \{a, a, b, c, c, c\}$ has cardinal six and not three as if it was a set of, say, ZFC (Blizard 1989). Then you could say that the vertices (and edges) are elements of a multiset and think that this could avoid the problem of their distinction. Yes, perhaps this could be done, but the mentioned philosophers should mention that, I mean, they should explain what they are presupposing as their background theory. In not doing that, we are free to suppose that they proceed as the scientist usually does, namely, to work with standard settings, that is, standard sets.

Consequently, as a result of considering the *standard* underlying logic (set theory), once nothing is said about that, we can surely assert that the ‘new category of objects’ introduced by these authors, namely, just *relational*s or just *weakly discernible* things cannot exist. Since this is a core point of this paper, let us insist: we should not be carelessness with the mathematical basis of our physical theory. If our quantum theory is grounded on one ‘classical’ system such as one of the above mentioned ones, ‘purely logical’ properties such as ‘to be identical with’ cannot be ignored, since they are part of this basis. This will be emphasised next.

3. On discerning kinds of properties

Another question posed by the mentioned authors concerns the distinction between what one could call ‘legitimate’ and ‘illegitimate’ properties to be used to make the

distinctions among quantum entities, or then that such properties would have a ‘physical content’ and so on. Other names could be used instead, and we shall try to be as simple as possible.

Suppose we have a (first-order or not) language L with a stock of unary predicate symbols whose arguments are individual terms (they are enough for our discussion here). Can we say that some of them are ‘physical’, ‘legitimate’, ‘yellow’ or ‘blue’? Let us consider just ‘yellow and ‘blue’ in our example in order to be neutral; later we go back to the physical case. This distinction aims at to discuss the proposal of some philosophers that the discerning properties in quantum physics would have ‘physical content’ and should not be ‘purely logical’, as the defined identity of the elements of the domain seems to be (the property ‘ I_a ’ discussed above).

The answer is that syntactically there is no way to distinguish among classes of predicate symbols. You can use many sorted languages or (equivalently, as we shall do now) suitable second-order predicates. But anyway you need to resort to semantics. Let us interpret L in a structure encompassing a non-empty domain D and so that the unary predicate symbols are associated to subsets of D . Hence we could, in principle, choose some of these subsets to stand for the yellow things and another subset for the blue things. Then, a collection of yellow things would be the extension of a ‘yellow predicate’ (a predicate holding just for yellow things), and the same would hold in the blue case. But this should be arbitrary if not motivated by some reason, that is, we should know in advance that some elements of D are yellow and that some are blue. In this sense, it is not the obedience to the predicates that make them yellow or blue but the other way around; the predicates come only to help the syntactic description of already known things: we would be begging the question. Furthermore, being elements of a *set*, even the elements of D of the same colour are discernible from one another and this is due to the force of the mathematics being used.

Philosophers such as Tomasz Bigaj say that we should not consider among the attributes of a thing properties such as “this very object” (Bigaj 2022, p. 36), which is equivalent to our I_a above, since the use of the symbol ‘ a ’ would report to some specific element. He continues (loc. cit.): “it is standard practice to exclude (...) [the] so-called *impure* properties, that is, properties that somehow involve reference to individual objects (and the relation of numerical identity).” Others such as Adam Caulton (and many others he mentions) would not agree with my proposed predicate I_a too; as Caulton says, “indiscernibility by monadic formulae [should not] contain individual constants or equality.” (Caulton 2022, p.579).

Why impose such a restriction? In the former case, I_a would be ‘impure’ in what sense? Why properties that involve reference to individual objects would be ‘impure’? A possible answer would be, in my opinion, the supposed fact that in using, say, the individual constant ‘ a ’, we need to suppose that the object a is already known, that is, it is identified from the start. The same could be said with respect to the identity rela-

tion, since in saying that something is ‘identical with something’, these ‘somethings’ must be given already. Another answer would be the desire for the maintenance of some weak form of PII (which I insist will be no more the very PII since some properties were ruled out of the domain of the universal quantifier) within a standard mathematical framework. These arguments, if cogent (the mentioned philosophers do not speak about their reasons), could be considered with care: in presupposing that the objects are given from the start, we are not necessarily committed with their identities; in supposing that there are eleven electrons in a Sodium neutral atom, we are not committed with the identity of the electrons, as seems quite evident. Even the reference to *some* particular electron, say that one in the outer shell of the Na atom does not entail that we are specifying a particular electron, but just some conditions (given by suitable quantum numbers) that need to be fulfilled in order we are authorised to say that there is an electron there. The reference (to the outer shell electron) is not to *that* electron specifically (say some called ‘Mike’), but to a condition to be satisfied by some quantum entity, that is, a physical law.

In a logical language, we can bypass Caulton’s and Bigaj’s questioning by defining I_a differently without making reference to a explicitly; by the way, let us recall that the standard language of ZFC doesn’t contain individual constants. Let x and y be individual variables (of the language of ZFC) and put $I_y(x) := x \in \{y\}$. Since the unitary set of any y can be get from the axioms, our definition is ready for the identity of y , whatever y can be. Furthermore, notice that the definition doesn’t use neither the identity symbol nor any individual constant, and is a legitimate monadic property since it has just one free variable.

But let us consider a little bit further this question as kinked to quantum physics. Really, this discussion is linked to the validity of the Principle of the Identity of Indiscernibles (PII) in quantum mechanics.¹⁰ In a second-order language, PII reads as follows:

$$(2) \quad \forall x \forall y (\forall X (X(x) \leftrightarrow X(y)) \rightarrow x = y),$$

where x and y are individual variables, X is a predicate variable and ‘ $=$ ’ is numerical identity. It is easy to see that if for every individual a , I_a is included as an attribute of a , then PII becomes trivial. Then philosophers provide ‘versions’ of PII by restricting the range of the universal quantifier, for instance assuming only monadic properties or also relational ones. But using our above language, let us restrict the range of the quantifier to yellow properties only, thus getting

$$(3) \quad \forall x \forall y (\forall X (Y(X) \rightarrow (X(x) \leftrightarrow X(y))) \rightarrow x = y).$$

where $Y(X)$ stands for ‘ X is the property ‘to be yellow’.

But in this case ‘ $=$ ’ cannot be numerical identity anymore, but some weaker relation of *indistinguishability relative to yellow properties*, since there could exist some

'other' property, not a yellow one, which could discern between the two objects; then, (3) is not PII. Really, PII (and numerical identity), as usually formulated, requires *all* properties; as Joseph Melia guessed, "if logician's 'V' doesn't mean 'all', what does it mean?" (Melia 1995).

So, there is no way to escape the conclusion that it is arbitrary to relegate some properties as 'impure' either in the discussions of PII or in looking for distinguishing properties. Logic is mandatory.

Of course what we said relative of yellow properties holds also for 'physical' properties, whatever they are. The mentioned philosophers claim that a physical property must be endowed with a 'physical' significance so that it can be measured, but this is unjustified without some interpretations, that is, justified in a purely syntactical setting. To say that a certain property has 'physical significance' makes no part of the theory's underlying logic except if we use a language as that described above in (3). Really, you should be able to choose a bunch of 'measurable properties', then represent them in the language and say that only these ones are able to provide any distinction among the involved objects. But, as in the case of indiscernibility according to yellow properties, this would be just indiscernibility relative to 'measurable' or 'physical' properties and not identity.

Furthermore, how could we justify that we cannot 'measure' whether an element of D does belong to $\{a\}$ or not? What is 'to measure'? Does it depend on the existence of a well-equipped laboratory full of PhDs? If we accept a wide definition of 'measurable' saying that a property P is measurable if it can be available in some context and under certain circumstances, say given by the theory's axioms, then we have all the reasons to say that the identity of a is a measurable property.

4. A last example

The carelessness with the underlying logic is not uncommon even among great mathematicians. Even Hermann Weyl made the same mistake in his discussion about the combinatorics of aggregates of individuals in the Appendix B of his celebrated book (Weyl 1949). In short, he was looking for what is important for quantum mechanics regarding such combinatorics, realizing that it is not the individuation of the entities that import, but their 'kinds' and quantities. In order to express that, he considered a set S with n elements endowed with an equivalence relation \sim . The equivalence classes C_1, \dots, C_k of the quotient set S/\sim were taken as the *states* the elements of S may be in, and the important thing for quantum physics is the cardinalities of these classes, so that the ordered "decomposition" $n_1 + \dots + n_k$ sums n . So, Weyl supposes, one is able to say that there are n_i elements in class C_i without commitment to the individuation of these entities.

Here goes Weyl's mistake. He cannot assert that the elements of the equivalence classes cannot be discerned from one another. As elements of a *set*, the considered entities are distinct elements, and the supposition that we can forget this and pay attention exclusively to the equivalence classes and their cardinals does not eliminate such a fact. Weyl's elements are individuals. For a discussion on this, see Krause (1991), and French & Krause (2006, §3.7).

5. Conclusions

To provide an end, let us consider again our language L , of first or second order. Can we say that there are available discerning formulas for any two objects we consider? Such formulas would be excluded from *symmetric* languages, that is, those that obey the following definition (Bigaj 2022, p.84): a language L is symmetric with respect to an intended interpretation \mathfrak{A} iff for every open formula $\phi(x_1, \dots, x_n)$ in L and any permutation $\sigma : D \rightarrow D$, $\mathfrak{A} \models \phi(x_1, \dots, x_n)$ iff $\mathfrak{A} \models \phi(\sigma(x_1), \dots, \sigma(x_n))$."

We should acknowledge that the language of ZFC is *not* symmetric since the whole universe of sets, taken as the structure \mathfrak{A} , has no nontrivial automorphisms (Jech 2003, p.66), and so there are open formulas that are not invariant by permutations, for instance, $x \in y$. But you can reply by suggesting that we should use ZFA instead. Due to the existence of permutation models, someone could argue that if we remain within a permutation model (or of another non-rigid structure), then there would be discerning formulas if they are restricted to the domain of the model. This is correct, but does not solve the problem; it is a theorem *about* ZFC (ZFA and the like) that *every structure can be extended to a rigid one* (da Costa & Rodrigues 2007), and so, even if some elements look like indiscernible things in the structure, they are seen to be discernible in the extended one.

Summing up, within 'standard' frameworks such as ZF, ZFA, etc., any object can always be discerned from any other at least by properties like its identity, which is a monadic property and there seems to be no indisputable way to say that some properties are yellow, physical, pure or illegitimate. In particular, within a standard framework encompassing classical logic and set theory, we cannot say that there are only relationals or only weakly discernible things. Standard logic forbids that.

References

Beth, E. 1966. *The Foundations of Mathematics: A Study in the Philosophy of Science*. New York: Harper & Row.
Bigaj, T. 2022. *Identity and Indiscernibility in Quantum Mechanics*. Cambridge: Palgrave-Macmillan.

Blizard, W. 1989. Multiset theory. *Notre Dame Journal of Formal Logic* 30(1): 36–66.

Bondy, J. A. & Murty, U. S. R. 2008. *Graph Theory*. Graduate Texts in Mathematics 244. New York: Springer.

de Barros, J. A.; Holik, F.; Krause, D. 2023. *Distinguishing indistinguishabilities: Differences Between Classical and Quantum Regimes*. Forthcoming by Synthese Library. Cham: Springer.

Bueno, O. 2010. A defense of second-order logic. *Axiomathes* 20: 365–383.

Caulton, A. 2022. Permutations. In E. Know; A. Wilson (ed.), *The Routledge Companion to Philosophy of Physics*, p.578–594. London: Routledge.

da Costa, N. C. A. & Rodrigues, A. A. M. 2007. Definability and invariance. *Studia Logica* 86(1): 1–30.

da Costa, N. C. A.; Krause, D.; Bueno, O. 2007. Paraconsistent logics and paraconsistency. In D. Jaquette (ed.), *Handbook of the Philosophy of Science*, Vol.5: Philosophy of Logic, p.791–912. Elsevier.

Dalla Chiara, M. L. & Toraldo di Francia, G. 1981. *Le Teorie Fisiche: Un'Analisi Formale*. Torino: Boringhieri.

Dalla Chiara, M. L.; Giuntini, R.; Krause, D. 1998. Quasiset theories for microobjects: a comparison. In E. Castellani (ed.), *Interpreting Bodies: Classical and Quantum Objects in Modern Physics*, p.142–152. Princeton: Princeton University Press.

Dalla Chiara, M. L.; Giuntini, R.; Greechie, R. 2004. *Reasoning in Quantum Theory: Sharp and Unsharp Quantum Logics*. Trends in Logic (Studia Logica Library) n.22. Dordrecht: Kluwer.

French, S. & Krause, D. 2006. *Identity in Physics: A Historical, Philosophical, and Formal Analysis*. Oxford: Oxford University Press.

Jech, T. 2003. *Set Theory: The Third Millennium Edition*. Revised and Expanded. Springer Monographs in Mathematics. Berlin and Heidelberg: Springer Verlag.

Krause, D. 1991. Multisets, quasi-sets and Weyl's aggregates. *Journal of Non-Classical Logic* 8(2): 9–39.

Krause, D. 2023. On identity, indiscernibility, and (non-)individuality in the quantum domain. *Philosophical Transactions of the Royal Society A* 381: 20220096. DOI 10.1098/rsta.2022.0096.

Krause, D. & Wajch, E. 2023. A Reappraisal of the Theory of Quasi-Sets with Emphasis on the Discussion about Quasi-Cardinals. *Preprint*.

Ladyman, J. & Ross, D. 2007. *Every Thing Must Go: Metaphysics Naturalized*. Oxford: Oxford University Press.

Melia, J. 1995. The Significance of Non-Standard Models. *Analysis* 55(3): 127–134.

Muller, F. A. 2011. The raise of relationals. *Mind* 124(493): 201–237.

Muller, F. A. & Saunders, S. 2008. Discerning fermions. *British Journal for the Philosophy of Science* 59: 499–548.

Ore, O. 1962. *Theory of Graphs*. Providence: American Mathematical Society.

Rodriguez-Pereyra, G. 2014. *Leibniz's Principle of Identity of Indiscernibles*. Oxford: Oxford University Press.

Russell, D. 1983. Anything goes. *Social Studies of Science* 13(3): 437–464.

Shapiro, S. 1991. *Foundations without Foundationalism: The Case for Second Order Logic*. Oxford Logic Guides, 17. New York: Oxford University Press.

Weyl, H. 1949. *Philosophy of Mathematics and Natural Science*. Princeton: Princeton University Press.

Wilson, R. J. 1996. *Introduction to Graph Theory*. 4th ed. Essex: Addison Wesley Longman.
Zee, A. 2023. *Quantum Field Theory, as Simply as Possible*. Princeton: Princeton University Press.

Notes

¹ My reading of this slogan does not go to some common interpretations which suppose that the slogan indicates that any theory (or logic) is as good as any other. In my opinion, the slogan is better read if indicates that the scientist is completely free to investigate. See Russell (1983).

² By ‘logic’ we mean here ‘great logic’ (*Logica Magna*); see Beth (1966, p.202), encompassing higher-order logic (at least second-order) or at least an amount of set theory, such as ZFC or ZFA. When just first-order logic is assumed, this is mentioned explicitly.

³ For a defense of the second-order approach, see Bueno (2010), Shapiro (1991).

⁴ In fact, the presence of contradictory sentences can be accepted without trivialisation only if the underlying logic is modified to some paraconsistent logic; see da Costa *et al.* 2007.

⁵ We are assuming von Neumann’s ordinals, so $\lambda < \nu$ means $\lambda \in \nu$; see (Jech 2003, p.19).

⁶ ZFC is the Zermelo-Fraenkel set theory with the Axiom of Choice, and ZFA is the Zermelo-Fraenkel set theory with the Axiom of Choice and atoms.

⁷ Philosophers use to distinguish between *synchronic* and *diachronic* identities. The first makes respect to an individual a at time t being the same (identical) to individual b at the same time t , while the second states that individual a at time t is identical with individual b at time t' .

⁸ ‘Properties’ are relations of arity one. They can be dealt with in the language of the theory either using primitive predicate symbols or as formulas with just one free variable.

⁹ We use the term ‘object’ as a neutral term to denote anything that can be conceptualised, in particular the ‘objects’ typical of quantum physics, such as electrons, protons, atoms and even molecules.

¹⁰ An important distinction is in order here. We should distinguish between the ‘logical’ PII, whose formulation is being presented here, and other forms of the principle such as those formulated for instance by Leibniz. We are discussing the logical version, and this ‘very (logical) PII’ requires the use of *all* properties and relations as we shall comment now.

Acknowledgments

I would like to thank the anonymous referee for several suggestions to improve the reading. The faults that remain are of course mine.