

Motor performance of elderly in a community in southern Brazil

Desempenho motor de idosos de uma comunidade do sul do Brasil

Susana Cararo Confortin¹
Aline Rodrigues Barbosa^{1,2}
Ana Lúcia Danielewicz²
Vandrize Meneghini³
Wagner Luiz Testa¹

Abstract – The objective of this study was to present reference values and assess the motor performance of the elderly in a community in the south of Brazil, according to sex and age group. Population-based household study, conducted with 477 elderly (≥ 60 years) from the municipality of Antônio Carlos, Santa Catarina state. Motor performance was verified by four tests: handgrip strength (KG), balance (four tasks), ‘chair stand’ and ‘pick up a pen’ (seconds). We use a score varying from 0 (unable) to 3 (good) in all the tests. We assessed 270 women with a mean age of 73.2 ± 8.82 years, and 207 men (73.3 ± 8.96 years). The results showed that the values (means, standard deviations and percentiles) were higher in men than those of the women in handgrip strength test and lower in the tests evaluated by time. The chi-square test showed that men had the highest prevalence of ‘good’ performance in the ‘chair stand’, balance and handgrip strength tests, when compared to the women ($p \leq 0.01$). With the advance of age there is a reduction in the prevalence ($p \leq 0.01$) of men and women with good performance and an increase in the prevalence of incapacity or poor performance in all the tests. The best motor performance is specific to the test, sex and age group. The men and the youngest age groups exhibit the best motor performance in all the tests.

Key words: Aging; Hand strength; Time and motion studies.

Resumo – O estudo teve como objetivo apresentar valores de referência e avaliar o desempenho motor de idosos de uma comunidade do sul do Brasil, de acordo com sexo e grupo etário. Estudo populacional, de base domiciliar, realizado com 477 idosos (≥ 60 anos) do município de Antônio Carlos, SC. O desempenho motor foi verificado por meio de quatro testes: força de prensão manual (Kg), equilíbrio estático (quatro tarefas), “sentar e levantar” e “pegar o lápis” (segundos). Foi usado escore de classificação para todos os testes que variou de 0 (incapaz) a 3 (bom). Foram avaliadas 270 mulheres ($73,2 \pm 8,82$ anos) e 207 homens ($73,3 \pm 8,96$ anos). Os resultados mostraram que os valores (médias, desvios padrão e percentis) dos homens, no teste de força de prensão manual, foram maiores do que os das mulheres e menores no testes avaliados por tempo. O teste qui-quadrado mostrou que os homens tiveram maior prevalência de desempenho “bom” nos testes “sentar e levantar”, equilíbrio e força de prensão manual, comparados às mulheres ($p \leq 0,01$). Com avanço da idade, houve redução na prevalência ($p \leq 0,01$) de homens e mulheres com desempenho bom e aumento na prevalência de incapacidade ou desempenho fraco, em todos os testes. O melhor desempenho motor é específico ao teste, sexo e grupo etário. Os homens e os grupos etários mais jovens apresentam melhor desempenho motor, em todos os testes realizados.

Palavras-chave: Envelhecimento, Estudos de tempo e movimento, Força de prensão manual.

1 Universidade Federal de Santa Catarina. Programa de Pós-graduação em Educação Física. Florianópolis, SC. Brasil

2 Universidade Federal de Santa Catarina. Programa de Pós-graduação em Nutrição. Florianópolis, SC. Brasil

3 Universidade Federal de Santa Catarina. Bacharelado Educação Física. Florianópolis, SC. Brasil

Received: 29 September 2012
Accepted: 02 January 2013



Licence
Creative Commons

INTRODUCTION

Motor performance tests are important objective measures in the assessment of the functional health of the elderly, as they allow the identification of physical/motor capacity involved in the performance diverse tasks related to daily living¹⁻³. The assessment of functional limitation through these types of tests can be indicative of the process of disabilities³, as it can predict cognitive deficit⁴, risk of fractures^{5,6}, hospitalization⁷, morbidities and mortality^{2,8}. In addition, the motor tests can characterize current health and assist in the delivery of intervention programs⁹.

In Brazil, there are few studies, with probabilistic samples and home collection that verify the motor performance of the elderly and present information regarding this performance in a way that provides a reference for the assessment of health. The 'SABE Survey'¹⁰ involved elderly residents in the urban area of the municipality of São Paulo. The study conducted by Pinheiro et al.¹¹ involved elderly residents in city with unfavourable socioeconomic conditions in north eastern Brazil. No studies involving elderly residents from rural areas, as in the south of the country, were found.

Brazil is a country which still has divergent socio-demographic, cultural and economic characteristics¹² and these characteristics may be reflected in living conditions, access to health care, health status and differences in the motor performance of the elderly^{2,9-11}. Therefore, it is necessary to conduct studies in various contexts, in order to have pertinent information from this contingent of the population. The objective of this study was to present reference values and assess the motor performance of the elderly from a community in the south of Brazil, where 71.5% of the elderly live in rural area, according to sex and age group¹³.

METHODS

This study, both population-and household-based, is part of the epidemiological research 'Saúde - AC', conducted in the municipality of Antônio Carlos, Santa Catarina (AC-SC). This municipality, 30 km from the capital of the state of Santa Catarina, in 2010, had a population of 7,458 inhabitants (12.8% were 60 or over), with 68.1% of the population residing in small, rural properties¹³. There is a Health Service Unit in the centre of the Antônio Carlos, in addition to three teams from the Family Health Strategy programme (FHS) which cover 100% of the municipality.

The study population was composed considering all the elderly registered, in 2009, on the FHS (n=917), attending to the age groups of 60-79 years and 80 years or more. For the individuals between 60 and 79 years (n=782) the calculated sample was of 471 individuals (margin of error of 5 percentile points, prevalence of 50% of unknown outcome, a test power of 80% and 15% sample loss). Random sampling was considered within each area of the FHS (area 1, n=175; area 2, n=140 and area 3, n=156). At the end of the collection (December 2010 to April 2011) 343 elderly were assessed (area 1, n=125; area 2, n=98 and; area 3, n=120) increasing the margin of error to 5.4

percentile points. Since the stratified sample was not proportional, sample weights were used for the analysis of data. The sample loss criteria were the following: absence of an adequate proxy respondent; individual absent from the municipality for a period longer than the field research or following three home visits; inability to access to the home due to the rural road conditions.

All the elderly (n=134) from the municipality that were 80 or over were assessed (February to April 2010). The elderly that were 80 years or over before 31st of May 2010 were placed in the 80 or over age group; the rest in the 60-79 age group.

We used a questionnaire based (short version) on the 'SABE Survey', a multicenter study undertaken in seven Latin American and Caribbean countries (http://hygeia.fsp.usp.br/sabe/Extras/Questionário_2000.pdf)¹⁴.

Data collection was conducted by undergraduate and postgraduate students (Physical Education and Nutrition) and by the research coordinator. The interviewers were trained prior to testing and refinement and calibration of the instrument (questionnaire and motor tests). The research coordinator was responsible for training and check the questionnaires.

The handgrip strength test (HGS) verified the motor performance of the superior limbs. The inferior limbs were evaluated by three tests related to the function of strength/resistance, mobility, balance and agility: 'chair stand'¹⁵, 'pick up the pen'¹⁶, assessed by time and balance¹⁵. Tests procedures were the same used in the 'SABE Survey'^{9,10} and will now be presented concisely.

Handgrip strength – assessed by a dynamometer (TAKEI), was taken using the arm that the subject considered dominant. The assessment was conducted twice with an interval of 1 minute and the highest value was recorded (kg). To assess the performance in the test, the values (kg) were distributed in percentiles, according to sex: unable = score 0 (unable); $\leq P_{25}$ = score 1 (poor); $> P_{25}$ a $\leq P_{75}$ = score 2 (average); $> P_{75}$ = score 3 (good).

The 'chair stand' test¹⁵ – the elderly began the test in the sitting position, arms crossed over the chest, after which they attempted to stand and sit five times, consecutively, as quick as possible, in ≤ 60 seconds. The performance in the test was verified by way of value distribution (seconds) in percentiles, according to sex: score 0 (unable); $\leq P_{25}$ = score 1 (poor); $> P_{25}$ a $\leq P_{75}$ = score 2 (average); $> P_{75}$ = score 3 (good).

The 'pick up the pen' test¹⁶ – the individual should crouch, pick up the pencil on the ground and return to the start position in ≤ 30 seconds. The performance in the test was verified by way of value distribution (seconds) in percentiles, according to sex: score 0 (unable); $\leq P_{25}$ = score 1 (poor); $> P_{25}$ a $\leq P_{75}$ = score 2 (average); $> P_{75}$ = score 3 (good).

Balance¹⁵ – verified by a test composed of 4 static tasks (10 seconds each), conducted with the individual standing, in sequence: (a) side-by-side stand; (b) full tandem stand; (c) maintain balance when standing only on the right leg (one-leg stand); (d) maintain balance when standing only on the left leg (one-leg stand). The score were the following^{9,10}: unable (0) = was not able to perform any of the 4 tasks; poor (1) = performed one task (side-by-side); average (2) = performed two tasks (side-by-side and full tandem stand); good (3) = successfully performed three or four tasks.

Before each test the interviewers explained and demonstrated the test and made sure that the task could be completed without any physical risk to the elderly. We take care in relation the footwear of individuals (tests related to the lower limbs), i.e., tests were performed with the individual barefoot or using any firm shoe.

The elderly unable of understanding the instructions due to a cognitive problem were excluded from the analyses. The elderly with physical limitations were included in the score tests as unable^{9,10}.

Information about family arrangements (living alone/accompanied), literacy (knows how to write and read - yes/no) and occupation throughout life (agriculture/other professions) were used to describe the sample.

For the descriptive analyses of the variables, measures, standard deviation, percentiles and proportion of individuals were used according to sex and age group (60-69, 70-79 and ≥ 80). The Kruskal-Wallis test verified the effect of the age group upon motor performance and the U test (Mann-Whitney) was used in the comparisons between men and women.

The distribution of the individuals, according to the motor performance test scores, sex and age group were shown in tables, using the chi-squared test. The exact Fisher test was used in the cases in which any frequency expected was less than five. The *Spearman* correlation was used to verify the relation between the test scores and the motor performance, according to sex.

All the analyses were weighted using the sample weight. The level of significance adopted was 5% ($\alpha = 0.05$). The data were doubly tabulated and analysed using the statistic programme SPSS[®] 16.0.

The ethics committee of the Universidade Federal de Santa Catarina (Protocol No. 189/09) approved the study.

RESULTS

The study involved 270 women (73.2 ± 8.82 years) and 207 men (73.3 ± 8.96 years), aged between 60 and 100 years. The majority of the elderly lived accompanied in the home (86.9%), worked in agriculture/farming throughout life (70.1%) and stated they were able to read and write (82.1%).

In the evaluation of motor performance, of the total 477 elderly analysed, 42 were unable to complete any test related to the performance of the inferior limbs and were included in the analyzes with score 0. Were also included with score 0, seven, 16 and 8 subjects who failed to perform the 'chair stand' test, 'pick up a pen' and balance tests, respectively. The test of handgrip strength was not realized by 17 elderly.

Table 1 presents the values from the HGS tests, 'chair stand' and 'pick up the pen', according to sex and age group. When compared to men, the women presented lower mean values in the HGS test ($p < 0.05$) and higher values in the tests assessed by time ($p < 0.05$), when compared with the men. With the increase in age, the men and women showed a significant reduction ($p < 0.001$) in mean values in the HGS and an increase in the time necessary to perform the 'pick up a pen' test ($p < 0.001$). The reductions occurred for the two extreme age groups (60-69 and ≥ 80 and over), except

for the handgrip test (men), whose differences were observed between younger age groups and the other age groups.

The data from Table 3 shows the distribution of the men and women, according to sex and performance in the tests. The male sex presented a higher prevalence of individuals with better results in the 'chair stand', balance and HGS tests, when compared with the female sex. The women showed a higher prevalence in incapability and poor performance in the 'chair stand' and balance tests, when compared with the men. The differences were significant between the sexes, in relation to performance in the 'chair stand' tests ($p \leq 0.001$), balance ($p \leq 0.001$) and HGS ($p \leq 0.021$).

Table 1. Means, standard deviation (SD) and medians, from the handgrip strength test (HGS), 'chair stand' and 'pick up a pen' according to sex and age group. Antônio Carlos, SC, Brazil, 2010/2011.

Tests	Age Group							
	60-69		70-79		≥ 80		All	
	Mean ± SD	Median	Mean ± SD	Median	Mean ± SD	Median	Mean ± SD	Median
Handgrip (kg)								
Female (263) *	22.4 ± 5.8	23.0 ^a	21.7 ± 5.8	21.0 ^b	19.0 ± 5.2	20.0 ^b	21.7 ± 5.8 [†]	22.0
Male (197) *	38.1 ± 8.8	40.0 ^a	29.2 ± 7.2	29.0 ^b	29.9 ± 7.0	31.0 ^b	33.5 ± 9.1 [†]	34.0
'Chair stand' (s)								
Female (226)	14.2 ± 6.9	13.0	14.2 ± 4.9	13.0	15.0 ± 5.7	13.5	14.3 ± 6.2 [†]	13.0
Male (182)	12.0 ± 4.6	11.0	12.8 ± 4.6	12.0	13.0 ± 4.6	12.0	12.4 ± 4.6 [†]	11.9
'Pick up a pen' (s)								
Female (233) *	2.2 ± 1.5	2.0 ^a	2.3 ± 1.9	2.0 ^a	3.4 ± 1.9	3.0 ^b	2.4 ± 1.7 [†]	2.0
Male (185) *	1.9 ± 1.2	2.0 ^a	2.0 ± 1.2	2.0 ^a	2.7 ± 1.3	2.0 ^b	2.0 ± 1.2 [†]	2.0

[†]Significant differences ($p < 0.05$) between the sexes (*Mann-Whitney*). * Values in reference to the comparison between the age groups (*Kruskal Wallis* test). a,b - values with different superscript letters were significantly different (*Mann-Whitney*).

Table 2. Percentiles of time spent conducting the 'chair stand' test (seconds), 'pick up a pen' test (seconds) and handgrip strength test (kg), according to sex and age group. Antônio Carlos, SC, Brazil, 2010/ 2011.

	Female				Male				All
	60-69	70-79	80	All	60-69	70-79	80	All	
Handgrip (n)	(115)	(72)	(76)	(263)	(78)	(65)	(54)	(197)	(460)
10	15.0	15.0	12.0	14.7	27.0	20.0	19.0	21.0	15.0
25	18.0	17.0	6.0	18.0*	34.2	24.0	25.8	28.0*	20.0
50	23.0	21.0	20.0	22.0	40.0	29.0	31.0	34.0	25.0
75	27.0	26.0	22.0	26.0*	44.0	33.7	35.0	41.0*	33.0
90	29.0	29.0	26.0	29.0	47.9	38.0	38.5	45.5	41.0
'Chair stand' (n)	(108)	(67)	(58)	(233)	(77)	(60)	(48)	(185)	(418)
10	8.0	9.0	9.0	9.0	8.0	7.0	8.0	8.0	8.0
25	10.0	12.0	11.0	11.0	9.0	10.0	10.0	10.0	10.0*
50	13.0	13.0	13.5	13.0	11.0	12.0	12.0	12.0	12.0
75	16.0	16.0	17.5	16.0	4.0	16.0	15.0	16.0	16.0*
90	19.0	19.0	23.8	20.0	18.0	19.0	19.0	19.0	19.0
'Pick up a pen' (n)	(110)	(64)	(52)	(226)	(77)	(60)	(45)	(182)	(408)
10	1.0	1.0	2.0	1.0	1.0	1.0	2.0	1.0	1.0
25	1.0	1.0	2.0	1.0	1.0	1.0	2.0	1.0	1.0*
50	2.0	2.0	3.0	2.0	2.0	2.0	2.0	2.0	2.0
75	2.0	3.0	4.0	3.0	2.0	2.0	3.0	2.0	3.0*
90	4.0	4.0	6.0	4.0	3.0	3.9	5.0	3.0	4.0

*Used to define the categories of performance.

The percentile distribution shows the values employed to determine the categories of motor performance (percentiles 25 and 75).

Table 3. Distribution (%) of the elderly according to sex and performance in the tests. Antônio Carlos, SC, Brazil, 2010/11.

	Unable	Poor	Average	Good	p*
'Chair stand'					
Female	10.7	19.0	50.9	19.4	≤0.001
Male	9.8	15.4	40.6	34.2	
Total	10.3	17.4	46.4	25.8	
'Pick up a pen'					
Female	8.7	2.8	22.4	66.1	0.465
Male	8.8	1.5	19.8	69.8	
Total	8.8	2.2	21.3	67.7	
Balance					
Female	9.3	19.8	29.5	41.4	≤0.001
Male	8.8	8.2	17.0	66.0	
Total	9.1	14.8	24.1	51.9	
Handgrip					
Female	1.2	15.5	46.4	37.0	≤ 0.021
Male	4.1	16.5	40.9	38.6	
Total	2.4	15.9	44.0	37.7	

*Chi-square Test.

With the advance of age, women and men present significant differences in performance in the four tests. Between the men, the prevalence of better performance reduced, in all the tests, with the advance of age. For the female sex, in the 'chair stand' test, the women aged 80+ exhibited better prevalence of good performance than those aged 70-79 (Table 4).

Table 4. Distribution (%) of the elderly according to age groups, sex and performance in the tests. Antônio Carlos, SC, Brazil, 2010/11.

Tests	Female			P	Male			P
	60-69	70-79	≥ 80		60-69	70-79	≥ 80	
Handgrip								
Unable	-	3.6	-	≤0,001*	28	7.2	-	≤0,001*
Poor	13.9	14.3	23.7		7.2	26.8	18.5	
Average	41.6	47.6	60.5		27.1	49.0	64.8	
Good	44.6	34.5	15.8		63.0	17.0	16.7	
'Chair stand'								
Unable	2.7	15.3	29.7	≤0.001*	4.9	13.2	18.2	≤0.007**
Poor	19.5	18.8	17.6		12.5	19.2	14.5	
Average	51.9	54.7	37.8		42.4	37.1	43.6	
Good	26.0	11.2	14.9		40.2	30.5	23.6	
'Pick up a pen'								
Unable	3.5	10.6	21.6	≤0.001*	4.9	13.2	11.1	≤0.001*
Poor	2.7	2.9	4.1		1.6	1.3	1.9	
Average	16.9	21.2	44.6		17.9	17.2	33.3	
Good	76.9	65.3	29.7		75.5	68.2	53.7	
Balance (n)								
Unable	4.5	12.4	18.4	≤0.001**	4.9	13.3	10.9	≤0.001*
Poor	19.8	18.2	23.7		8.2	8.7	7.3	
Average	18.7	43.5	36.8		10.9	20.7	27.3	
Good	57.1	25.9	21.1		76.1	57.3	54.5	

* Fisher's exact test. ** Chi-square Test.

The scores of the tests were positively correlated with statistical significance ($p \leq 0.01$) (Table 5).

Table 5. Spearman correlation coefficient for testing motor performance in elderly. Antônio Carlos, SC, Brazil, 2010/2011.

	r_{spearman}	r_{spearman}	r_{spearman}
	'Chair stand'	'Pick up a pen'	Balance
Female			
HGS	0.344*	0.253*	0.330*
"Chair stand"		0.484*	0.506*
"Pick up a pen"			0.348*
Male			
Handgrip	0.454*	0.426*	0.383*
"Chair stand"		0.509*	0.409*
"Pick up a pen"			0.508*

* $p \leq 0.01$

DISCUSSION

According to our knowledge, this is the first home-based population study in Brazil that presents reference values and assesses the motor performance of elderly in rural and urban areas. The previous studies covered urban areas only^{10,11}.

The results showed that the men exhibited superior values to the women in the HGS, completed the tests in less time and had a higher prevalence in good performance in all the tests, as is described in the literature⁹⁻¹¹. The women exhibited the higher percentage of incapability and/or poor performance in the tests. With advancing age, male and female showed a reduction in the proportion of individuals with better results in all tests.

The differences between the sexes in motor performance may be explained by the hormonal characteristics and their effects on body composition¹⁷. The women exhibit a higher quantity of body fat while the men showed higher levels of testosterone and a higher quantity of muscle mass. The women are more affected by chronic diseases that cause pain and limit joint movements¹⁸ and exhibit a higher prevalence of excess weight¹⁰, factors that hinder mobility¹⁹ and balance^{2,20}. It is believed that the lifestyle of men, working in farming, could have contributed to the better performance in the tests, even at more advanced ages.

Comparing the results of the present study with other studies that used the same test protocols and instruments, can be verified that men and women from AC-SC showed higher values of HGS, compared to the elderly from São Paulo¹⁰, Barbados and Cuba⁹. In São Paulo were evaluated 1894 elderly aged 60 years or more, and in Barbados and Cuba⁹, 1508 and 1905 elderly were evaluated, respectively. The men from AC-SC, were faster than those of Cuba⁹ and of São Paulo¹⁰ in the 'chair stand' test and slower than the elderly of São Paulo¹⁰ in the 'pick up a pen' test. The women

from AC-SC completed the 'chair stand' test in an inferior time than the women from Cuba⁹. In the 'pick up a pen' test the women from this study were faster than the women from Cuba, Barbados⁹ and São Paulo¹⁰.

In relation to the percentile distribution, the values used to define the categories of performance were equal to those observed in the 'SABE Survey' São Paulo¹⁰ for the 'chair stand' test and to those from Barbados⁹ and Lafaiete Coutinho-BA¹¹ in the 'pick up a pen' test. In the HGS the elderly from AC-SC exhibited higher values than those of the elderly from São Paulo¹⁰, Lafaiete Coutinho-BA¹¹, Barbados and Cuba⁹. However, it is worth noting that the study by Pinheiro et al.¹¹ used a different dynamometer to that used in AC-SC, São Paulo¹⁰, Barbados and Cuba⁹, which may interfere in the identified values.

The comparison with other populations, even using the same methodology and instruments should be viewed with caution. The differences may be due to differences in the criteria used for the sample selection and/or exclusion of elderly, number of participants, the better ability of individuals, ethnic differences, environmental influences¹⁰ and willingness to participate, besides differences in nutritional status and physical activity level.

The results showed that the prevalence of individuals with better performance was specific to each test, varying between sex and age group. However, with the advance of age, men and women showed a decrease in the prevalence of good performance and the increase in the percentage of those unable or with poor performance as verified in other studies^{9,10,21}.

The highest prevalence of chronic disease, the reduction of physical activity, in addition to the normal physiological alterations of aging (reducing of muscle mass, the decline in the levels of determined hormones¹⁷, cognitive decline²², alterations in the sensory systems²³ and the nutritional status²⁰, among others), may explain the motor decline with the advance of age.

The correlations between motor performance scores for both sexes were positive and significant between all tests, suggesting that elderly that exhibit good performance in the first task tend to present similar performance in the second. These findings are similar to those verified in the 'SABE Survey'/São Paulo¹⁰ and the study conducted in Lafaiete Coutinho-BA¹¹.

The cross-sectional design does not allow to verify the effects of aging on the reduction of motor performance and / or increase in functional limitation. However, the results are consistent with findings from other studies. The results allow the identification of the prevalence of elderly with poor performance/functional limitation and, as the results were presented to the municipality of AS-SC (Secretary of Health and Social Assistance), they could contribute to the planning of actions to improve the health of the elderly. The study included a representative sample of the elderly population of the municipality, guaranteeing internal validation. The other strong point of the study is the possibility of presenting reference values for a set of motor tests that could be used as a reference for elderly from municipalities in the south of Brazil, with similar characteristics.

CONCLUSION

Data from this study showed that the men and the youngest elderly exhibited better performance in the motor tests, compared to the women and the eldest individuals, respectively. The highest prevalence of incapability and best motor performance is specific to the test, sex and age group.

Acknowledgments

The authors want to thank Antonio Carlos' Board of Health and Social Assistance, the Health Community Agents, and the oldest old attendees on this research. Our thanks also go to the National Council of Technological and Scientific Development (CNPq- Process 478073/2009-7), for financing the project. Meneghini V. received a scholarship grant from National Council of Technological and Scientific Development (PIBIC); Confortin S.C. and Danielewicz A.L. received a master's scholarship grant from Coordination of Improvement of Higher Education in Brazil (Capes-Reuni).

REFERENCES

1. Guralnick JM, Ferruci L. Assessing the building blocks of function utilizing measures of functional limitation. *Am J Prev Med* 2003;25(3 Suppl):112-21.
2. Castaneda-Sceppa C, Price LL, Noel SE, Midle JB, Falcon LM, Tucker KL. Physical function and health status in aging Puerto Rican adults: the Boston Puerto Rican health study. *J Aging Health* 2010;22(5):653-72.
3. Seidel D, Brayne C, Jagger C. Limitations in physical functioning among older people as a predictor of subsequent disability in instrumental activities of daily living. *Age Ageing* 2011;40(4):463-9.
4. Wang L, Larson EB, Bowen JD, van Belle G. Performance-based physical function and future dementia in older people. *Arch Intern Med* 2006;166(10):1115-20.
5. Piirtola M, Vahlberg T, Isoaho R, Aarnio P, Kivela SL. Predictors of fractures among the aged: a population-based study with 12-year follow-up in a Finnish municipality. *Aging Clin Exp Res* 2008;20(3):242-52.
6. Ensrud KE, Ewing SK, Cawthon PM, Fink HA, Taylor BC, Cauley JA, et al. A comparison of frailty indexes for the prediction of falls, disability, fractures, and mortality in older men. *J Am Geriatr Soc* 2009;57(3):492-8.
7. Cawthon PM, Fox KM, Gandra SR, Delmonico MG, Chiou C, Anthony MS, et al. Do muscle mass, muscle density, strength and physical function similarly influence risk of hospitalization in older adults? *J Am Geriatr Soc* 2009; 57(8):1411-9.
8. Silventoinen K, Magnusson PK, Tynelius P, Batty GD, Rasmussen F. Association of body size and muscle strength with incidence of coronary heart disease and cerebrovascular diseases: a population-based cohort study of one million Swedish men. *Int J Epidemiol* 2009;38(1):110-8.
9. Barbosa AR, Miranda LM, Guimarães AV, Corseuil HX, Corseuil MW. Age and gender differences regarding physical performance in the elderly from Barbados and Cuba. *Rev Saude Publica* 2011;13(1):54-66.
10. Barbosa AR, Souza JMP, Lebrão ML, Marucci MFN. Functional limitations of Brazilian elderly by age and gender differences: data from SABE Survey. *Cad Saúde Pública* 2005;21(4):1177-85.
11. Coqueiro RS, Fernandes MH, Barbosa AR. Motor performance of the elderly in northeast Brazil: differences with age and sex. *Rev Esc Enferm USP* 2013; 47(1):125-33.

12. Massuquetti A, Franco Junior MCR. O sul e o Nordeste no Brasil: Uma análise das diferenças no desenvolvimento socioeconômico destas regiões. II Encontro de Economia Catarinense, Chapecó, SC, 2008. Available from: <http://www.apec.unesc.net/II%20EEC/sessoes_tematicas/Especiais/Artigo7.pdf> [2012 Set 15].
13. Instituto Brasileiro de Geografia e Estatística (IBGE). Censo 2010. Available from: <<http://www.ibge.gov.br/censo2010/>> [2012 Jun 20].
14. Lebrão ML, Laurenti R. Saúde, bem-estar e envelhecimento: o estudo SABE no município de São Paulo. *Rev Bras Epidemiol* 2005;8(2):127-41.
15. Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol Med Sci* 1994;49(2):M85-M94.
16. Reuben DB, Siu AL. An objective measure of physical function of elderly outpatients – The Physical Performance Test. *J Am Geriatr Soc* 1990;38:1105-12.
17. Valentine RJ, Misic MM, Rosengen KS, Woods JA, Evans EM. Sex impacts the relation between body composition and physical function in older adults. *Menopause* 2012;16(3):518–23.
18. Kim IH, Chun H, Kwon JW. Gender differences in the effect of obesity on chronic diseases among the elderly Koreans. *J Korean Med Sci* 2011;26(2):250-7.
19. Koster A, Penninx BWJH, Newman AB, Visser M, van Gool CH, Harris TB, et al. Lifestyle factors and incident mobility limitation in obese and non-obese older adults. *Obesity* 2007;15(12):3122-32.
20. Barbosa AR, Souza JMP, Lebrão ML, Marucci MDFN. Estado nutricional e desempenho motor de idosos de São Paulo. *Rev Assoc Med Bras* 2007;53(1):75-9.
21. Schlüssel MM, Anjos LA, Vasconcellos MTL, Kac G. Referente values of hand-grip dynamometry of health adults: A populatin-based study. *Clin Nutr* 2008; 27(4):601-7.
22. Kato-Narita EM, Nitrini R, Radanovic M. Assessment of balance in mild and moderate stages of Alzheimer's disease. *Arq Neuropsiquiatr* 2011;69(2-A):202-7.
23. Toledo DR, Barela JA. Diferenças sensoriais e motoras entre jovens e idosos: contribuição somatossensorial no controle postural. *Rev Bras Fisioter* 2010; 14(3):267-75.

Corresponding author

Aline Rodrigues Barbosa
Departamento de Educação Física –
Centro de Desportos
Universidade Federal de Santa
Catarina - Campus Trindade,
88940 – Florianópolis, SC, Brasil
E-mail: aline.r.barbosa@ufsc.br
alinerb13@yahoo.com.br