

**Artigo original**

Betty M. Pérez¹
Maura Vásquez²
Maritza Landaeta-Jiménez³
Guillermo Ramírez²
Coromoto Macías-Tomei⁴

ANTHROPOMETRIC CHARACTERISTICS OF YOUNG VENEZUELAN SWIMMERS BY BIOLOGICAL MATURITY STATUS

CARACTERÍSTICAS ANTROPOMÉTRICAS DE JOVENS NADADORES VENEZUELANOS POR ESTÁGIO DE MATURAÇÃO BIOLÓGICA

RESUMO

Existe limitada informação sobre a maturação de crianças e adolescentes atletas, uma vez que a maturação é um fator interveniente na performance. O presente estudo mensurou variáveis antropométricas de 115 nadadores Venezuelanos (idade 7-17 anos), em diferentes estágios de maturação biológica. Os procedimentos antropométricos usados foram os recomendados pela ISAK. Os estágios de maturação genital de Tanner (G-5) foram utilizados para subdividir a amostra nas três categorias sugeridas por Nicoletti: pré-puberal (G1), puberal inicial (G2-3) e puberal avançada (G4-5). Foi utilizada a análise discriminante passo a passo (stepwise), tendo como critério de seleção o aumento da porcentagem de indivíduos classificados corretamente. Foram utilizadas 35 variáveis, das quais foram selecionadas: estatura, dobras cutâneas tríceps e subescapular, perímetros da perna e do braço relaxado, massa corporal, e os comprimentos do antebraço e coxa. Essa análise gerou um novo grupo composto por 87% do grupo original. Duas funções lineares discriminantes (LDF1 e LDF2) emergiram: LDF1 diferenciou os indivíduos pré-púberes dos púberes avançados, tendo incluído as variáveis: estatura, massa corporal, comprimento do antebraço e coxa, e os perímetros do braço e da perna. A segunda função, a LDF2, diferenciou os indivíduos no início da puberdade dos demais grupos. O segundo grupo de variáveis incluiu: a dobra cutânea do tríceps e o perímetro da perna. Os intervalos de confiança demonstraram que duas variáveis de tamanho, duas de gordura subcutânea, duas de robustez esquelética e duas de muscularidade relativa são suficientes para guiar os técnicos, na classificação de jovens nadadores, em categorias que melhor aproximem do seu estágio maturacional. Somente as dobras cutâneas do tríceps e subescapular mostraram superposição dos intervalos de confiança nas três categorias.

Palavras-chave: adolescente, maturação sexual, nadador, cineantropometria, análise discriminante.

ABSTRACT

There is limited information available on the maturation of child and adolescent athletes, considering that maturation is a factor that can affect performance. This study evaluates the anthropometry of 115 male Venezuelan swimmers (aged 7-17), at differing stages of biological maturity. The procedures employed adhere to International Society for the Advancement of Kinanthropometry guidelines. Tanner's stages of genital maturation (G1-5) were used to subdivide the sample into the three categories posited by Nicoletti, prepuberty (G1), initial puberty (G2-3) and advanced puberty (G4-5). Forward stepwise discriminant analysis was performed, using increase in hit rate as the selection criterion. A total of 35 variables were evaluated, from which stretch stature, triceps skinfold, subscapular skinfold, calf girth, relaxed arm girth, body mass, forearm length and thigh length were selected. This set produced a high aggregate hit rate (87%) of original grouped cases. Two linear discriminant functions (LDF1 and LDF2) emerged: LDF1 is capable of discriminating between prepuberty and advanced puberty and consists of stretch stature, body mass, forearm length, thigh length, relaxed arm circumference and calf circumference. LDF2 differentiates initial puberty from the other groups. This second set of discriminant variables comprises triceps skinfold and calf circumference. Confidence intervals demonstrate that two variables for body size, two for subcutaneous fatness, two for skeletal robustness and two for relative muscularity suffice to guide coaches when placing young swimmers into categories that best approximate the variability of maturation. Only triceps and subscapular skinfolds exhibited overlapping intervals in the three categories.

Key words: adolescents, sexual maturation, swimmers, kinanthropometry, discriminant analysis.

¹ Institute of Economical and Social Research. FaCES/ Central University of Venezuela.

² School of Statistics. FaCES/ Central University of Venezuela.

³ Bengoa Foundation.

⁴ Simón Bolívar University.

INTRODUCTION

The number of children and youths participating in high-level competitive sports is on the rise, and it is a fact that during prepubescence, and especially at maturation, many somatic, functional and physiological changes occur which increase the biological variability among children of the same chronological age. (1), (2) Despite this, the majority of studies focus on chronological age groups, even though classification on the basis of chronological age, needless to say, is of limited utility as an indicator of maturity status during this period of life. (3) On the other hand, according to several authors, different levels of performance within a given age group are often the result of the variations observed in the timing and tempo of these changes, which produce different strength, speed and endurance outcomes. (4)

Many studies have assessed the physical characteristics of athletic populations at different levels of competition, from Olympic to national level. The results of these studies suggest that a particular profile is success associated with performance. (5) On the other hand, data for Latin American child and adolescent swimmers are less often considered and hence information is limited. What is more, although maturation status may contribute to elite performance, and, in the case of swimming, the bodies of adolescent participants in this sport are still in the developmental phase, very little information is provided on the physical structure of young Latin American swimmers. However, the assessment of sexual maturation is no easy task, especially not in the presence of limitations that jeopardize a proper physical assessment. Accordingly, the present study was directed at the formulation of a classification rule, based on a set of anthropometric variables, to produce an approximation of maturation categories that could serve as a guide for coaches.

METHODS

Subjects

The sample analyzed here was part of a project on the growth and nutritional status of young Venezuelan swimmers, a larger collaborative research project carried out by the Central University of Venezuela and Fundacredesa and entitled, "Assessment of Growth, Nutritional Status, Food Intake and Socioeconomic Conditions of Young Venezuelan Swimmers". (6). The athletes who volunteered to enroll on the study were members of the Miranda State team, which currently holds the national swimming championship. Data were collected mid-season. Younger swimmers trained in one two-hour session a day, five days a week, while older swimmers trained in daily two-hour sessions, six days a week. About 60% of the athletes sampled

competed in regional and/or national meets and a smaller group was already participating in international events. Testing procedures were fully explained and informed consent was obtained from parents in writing. We selected for consideration just one hundred and fifteen male child and adolescents swimmers (aged 7-17) from the total cross-sectional sample of the whole project, as meeting the established criteria based on a complete set of anthropometric variables and the assessment of sexual maturation.

Thirty-five anthropometric variables were selected for the study, and their values measured in accordance with the ISAK guidelines. (7) Data was collected for all participants on the following anthropometric variables. There were four body size variables, these being stretch stature, body mass, sitting height and arm span, and eight subcutaneous fatness variables, namely triceps, subscapular, biceps, iliac crest, supraspinal, abdominal, front thigh and medial calf skinfolds. Indicators of skeletal robustness comprised seven length measurements, acromiale-radiale (arm length), radiale-styilion (forearm length), midstyilion-dactyilion (hand length), trochanterion-tibiale laterale, tibiale laterale, tibiale mediale-spyrion tibiale and foot and seven breadth measurements, biacromial, biiliocristal, transverse chest, anterior-posterior chest depth, biepicondilar humerus, wrist and biepicondilar femur. There were nine variables indicative of relative muscularity, all girths, specifically relaxed arm, arm flexed and tensed, forearm, wrist, chest, waist, gluteal (hip), thigh and calf circumferences.

Stretch stature, sitting height and arm span were determined to the nearest 0.1 cm. and body mass to the nearest 0.1 kg., employing a portable stadiometer, a wall scale Brocca plane and a portable scale, respectively. Skinfold thicknesses were measured using SlimGuide calipers (10 g/mm² constant pressure), a widely-used type of plastic caliper that gives results similar to the Harpenden (8). These dimensions were taken in duplicate, and the means of the trial values entered on the anthropometric datasheet. If a third trial was required, the median of the three measures was taken as the representative value. Direct and projected lengths were measured to the nearest 0.1 cm using an anthropometer and a segmometer respectively. Breadths and circumferences were also recorded to the nearest 0.1 cm with an anthropometer and a Lufkin steel tape, respectively.

Measurements were taken by one uncertified anthropometrist, and one Level 1 and one Level 2 ISAK certified anthropometrist, all supervised by an ISAK Level 3 Instructor (BP), who also measured many of the subjects. Prior to the project all measurers were found to meet the ISAK criteria for technical errors of measurement of <5% for skinfolds and <1% for all other variables. (9), (10)

The sample was subdivided on the basis of

Tanner's (11) sexual maturation stages of genitalia (G1-5) into the three categories posited by Nicoletti: Prepuberty (G1), Initial puberty (G2-3) and Advanced puberty (G4-5). (12) Assessment of sexual maturation was based on clinical examination. Maturity rate was monitored by a physician (CM-T). The development of secondary sex characteristics such as development of the genitalia has been summarized in the five-stage scales of the Marshall-Tanner method. (13) Therefore, groups were partitioned into the categories of prepuberty (stage 1), characterized for the appearance of male genitalia as in early childhood; initial puberty (stages 2 and 3) for initial and continued development of those organs, and late puberty (stages 4 and 5) as indicative of their continued and adult or terminal development.

Statistical analysis

In order to build a rule for classification based on the identification of anthropometric variables affording greater power for predictive purposes among prepubescents, initial pubescents and advanced pubescents, a forward stepwise discriminant analysis was performed, using increase in hit rate as the selection criterion. (14) The best subset of indicators was selected according to their relative contribution to the proportion of correct classification, both globally and within groups. 95% confidence intervals were estimated for means of the selected variables. Statistical analyses were performed using SPSS software (version 10.0) and SPAD 4.5, with significance set at $p < 0.05$.

RESULTS

Mean anthropometric variables for each maturity group are summarized in Table 1. In general, for the majority of the variables, mean values increase from prepubescence to advanced pubescence as expected. The exception was skinfold measurements, which showed a tendency to diminish. Subscapular skinfolds, however, behaved in line with the other dimensions.

There was a total number of 35 variables, from which stretch stature, triceps and subscapular skinfold, calf and relaxed arm girths, body mass, and forearm and thigh lengths were selected as the best subset of anthropometric variables, in terms of their predictive capabilities. Ranking by stepwise selection is presented in Table 2.

This set of variables produced a high aggregate apparent hit rate of 87%. The hit rate increases through the maturity groups from 80% in prepuberty to 93% in advanced puberty. During the selection process it was observed that stretch stature, triceps and subscapular skinfolds accounted for 75% of correctly classified prepubescents. When calf circumference, relaxed arm and body mass were included the hit rate reached 80%. All variables except lengths were required to reach

83% correct classification of initial pubescents. Finally, the most powerful variables for discriminating advanced puberty were stretch stature and triceps skinfold, which identified 89%.

Two linear discriminant functions (LDF1 and LDF2) capable of separating sexual maturation groups emerged when Wilks' lambda test ($p=0.000$) was applied. These functions were able to account for a substantial proportion of the differences between groups, as indicated by their squared canonical correlations (0.66 and 0.39 respectively). The resulting correlation structure is shown in Table 3.

These results indicate that the scores from LDF1 are basically made up of stretch stature, body mass, forearm length, thigh length, relaxed arm circumference and calf circumference measurements. LDF1 may account for separation of prepuberty and advanced puberty due to important differences in the discriminant variables. The LDF2-associated separation of initial puberty from the other groups may be attributed to the adiposity level in the triceps skinfold (Figure 1).




Figure 1. Canonical Discriminant Functions LDF Plot.

Table 4 summarizes the 95% confidence intervals for means of the eight variables identified in the discriminant analysis, by prepubescents, initial pubescents and advanced pubescents. According to these results, two variables for body size, two for subcutaneous fatness, two for skeletal robustness and two for relative muscularity suffice to guide coaches when placing young swimmers into the categories that best approximate the variability of maturation. Only triceps and subscapular skinfolds present overlapping subscapular intervals in the three categories, which is characteristic of the skinfolds in these athletes.

DISCUSSION

Despite the enormous amount of descriptive material available on the physiques of athletes, data concerning maturation in young Venezuelan competitive swimmers have not been collected to the

Table 1. Anthropometric variables (mean \pm standard deviation) by maturity status

Variables	Prepuberty (n=20)	Initial prepuberty (n=53)	Advanced prepuberty (n=27)
Body Size			
Stretch stature (cm)	136.4 \pm 6.7	152.6 \pm 10.8	170.0 \pm 7.9
Body mass (Kg)	31.6 \pm 5.1	44.2 \pm 9.8	58.5 \pm 10.3
Sitting height (cm)	71.3 \pm 2.6	77.8 \pm 6.4	86.8 \pm 4.1
Arm span (cm)	137.7 \pm 7.8	156.6 \pm 13.1	174.7 \pm 10.9
Subcutaneous Fatness			
Triceps skinfold (mm)	8.1 \pm 3.3	9.2 \pm 2.7	7.0 \pm 2.5
Subscapular skinfold (mm)	6.1 \pm 2.2	6.3 \pm 1.4	7.1 \pm 1.6
Biceps skinfold (mm)	5.1 \pm 2.3	5.2 \pm 1.7	3.7 \pm 0.7
Iliac crest skinfold (mm)	9.3 \pm 5.0	10.2 \pm 3.8	9.4 \pm 3.7
Supraspinal skinfold (mm)	5.8 \pm 3.2	6.2 \pm 2.4	6.0 \pm 1.9
Abdominal skinfold (mm)	8.3 \pm 4.3	9.6 \pm 3.8	8.9 \pm 3.3
Front thigh skinfold (mm)	12.2 \pm 3.6	13.5 \pm 4.3	10.4 \pm 2.4
Medial calf skinfold (mm)	8.7 \pm 3.0	10.3 \pm 3.7	8.2 \pm 1.9
Skeletal Robustness			
Lengths			
Arm length (cm)	25.5 \pm 1.8	28.7 \pm 2.3	31.9 \pm 2.1
Forearm length (cm)	20.0 \pm 1.3	23.3 \pm 2.2	26.1 \pm 2.5
Hand length (cm)	15.6 \pm 1.0	17.7 \pm 1.5	19.1 \pm 1.0
Trochanterion-tibiale laterale length (cm)	33.3 \pm 3.5	36.9 \pm 3.4	41.7 \pm 3.3
Tibiale laterale length (cm)	40.9 \pm 7.4	43.1 \pm 3.8	46.8 \pm 2.5
Tibiale mediale-spyrion tibiale length (cm)	30.5 \pm 2.1	34.4 \pm 2.9	38.6 \pm 2.6
Foot length (cm)	21.5 \pm 1.6	23.8 \pm 1.7	25.9 \pm 1.4
Breadths			
Biacromial breadth (cm)	30.0 \pm 1.9	33.3 \pm 3.4	38.4 \pm 2.6
Biliocrystal breadth (cm)	19.6 \pm 1.9	22.5 \pm 2.4	24.8 \pm 2.8
Transverse chest breadth (cm)	20.4 \pm 1.5	23.3 \pm 2.6	26.0 \pm 1.9
Anterior-posterior chest depth (cm)	14.0 \pm 1.3	16.0 \pm 1.9	18.2 \pm 1.8
Biepicondilar humerus breadth (cm)	5.5 \pm 0.4	6.2 \pm 0.4	6.8 \pm 0.4
Wrist breadth (cm)	4.4 \pm 0.4	4.9 \pm 0.4	5.4 \pm 0.4
Biepicondilar femur breadth (cm)	8.4 \pm 0.5	9.2 \pm 0.6	9.6 \pm 0.5
Relative Muscularity			
Relaxed arm (cm)	20.0 \pm 2.2	23.3 \pm 2.7	26.8 \pm 2.9
Arm flexed and tensed (cm)	21.3 \pm 2.3	24.7 \pm 2.9	28.8 \pm 3.1
Forearm (cm)	19.5 \pm 1.4	22.0 \pm 2.0	24.5 \pm 2.1
Wrist (cm)	13.1 \pm 0.6	14.6 \pm 1.2	15.8 \pm 0.9
Chest (cm)	67.6 \pm 4.6	76.4 \pm 7.5	87.1 \pm 6.7
Waist (cm)	58.4 \pm 3.6	64.8 \pm 5.0	71.0 \pm 6.4
Gluteal (hip) (cm)	67.6 \pm 4.4	77.2 \pm 7.3	85.0 \pm 5.8
Thigh (cm)	40.2 \pm 3.4	46.3 \pm 4.8	49.8 \pm 4.2
Calf (cm)	26.7 \pm 2.0	30.5 \pm 2.9	

Table 2. Stepwise ordering of anthropometric variables by prediction capability

Variables	Final rank	Step						
		1	2	3	4	5	6	7
stretch stature	1	1	1	1	1	1	1	1
triceps	2		2	2	2	2	2	2
subscapular	3			3	3	3	3	3
Calf circumf.	4				4	4	4	4
relaxed arm	5					5	5	5
body mass	6						6	6
forearm length	7							7
thigh length	8							
%Clas_Global	87	72	76	78	78	79	83	83
%Clas_Prepub	80	65	70	75	75	75	80	80
%Clas_InitPub	87	72	72	76	76	79	83	79
%Clas_AdvPub	93	78	89	85	85	82	85	93

Table 3. Discriminant functions (LDF1, LDF2)

Variable	Structure Matrix	
	Function	
	LDF1	LDF2
Stretch stature	0.898**	0.045
Body mass	0.737**	0.009
Forearm length	0.706**	0.147
Relaxed arm	0.643**	0.034
Thigh length	0.636**	-0.056
Calf circumference	0.609**	0.214*
Subscapular skinfold	0.167*	-0.105
Triceps skinfold	-0.133	0.340**
Canonical Correlation	0.810**	0.628
Wilk's Lambda	0.208**	0.605*

Pooled within-groups correlations between discriminating variables and standardized canonical discriminant functions.

Variables ordered by absolute size of correlation within function

**Significant at the 0.01 level

*Significant at the 0.05 level

Table 4. 95% confidence interval for means of anthropometric characteristics that discriminate maturity status

Variables	Prepubescent(<i>n</i> =20)		Initial pubescent(<i>n</i> =53)		Advanced pubescent (<i>n</i> =27)	
	lower bound	upper bound	lower bound	upper bound	lower bound	upper bound
Body Size						
Body mass (Kg.)	29,2	34,0	41,5	46,9	54,4	62,6
Stretch stature (cm.)	133,2	139,5	149,6	155,5	166,8	173,1
Subcutaneous Fatness						
Triceps skinfold (mm.)	6,6	9,7	8,4	9,9	6,1	8,0
Subscapular skinfolds (mm.)	5,1	7,2	5,9	6,7	6,5	7,8
Skeletal Robustness						
<i>Lengths</i>						
Forearm length (cm.)	19,0	21,1	22,67	24,0	25,7	28,0
Thigh length (cm)	31,6	34,9	35,9	37,8	40,4	43,0
Relative Muscularity						
Relaxed arm (cm.)	19,0	21,1	22,6	24,0	25,7	28,0
Calf (cm.)	25,7	27,6	29,7	31,3	32,2	33,9

requisite level of detail. The present study showed that a reduced number of variables could be taken into account for the classification of swimmers within the maturation process. Changes in the relationships between the variables over time affect the weight that each variable brings to the discriminant analysis.

Based on the results above, eight measurements could be used to classify these swimmers in groups reflecting maturation categories. As elicited in Table 2, stretch stature is by itself the most predictive capability variable. Differences in stature are in line with the biological variability of growth among maturation groups. Stature is the most commonly used indicator of somatic maturation, as it marks an onset of physical development during puberty. Most other indicators, such as changes in body dimensions, proportions and composition, follow the pace set by this variable. Swimmers generally exhibit greater height than the average population, in conjunction with earlier maturation. (15) This fact was particularly true among initial and advanced pubescent, and is in agreement with the tendency in the general population of high-income boys and girls toward earlier maturation that has already been observed in the Caracas longitudinal study. (16) Furthermore, based on previous studies, (17) it will be noted that Venezuelan swimmers were taller and heavier in comparison with Cuban and Mexican swimmers.

Additionally in this group of swimmers it is important to note that an important marginal increase in apparent hit rate when triceps and subscapular skinfolds are taken in conjunction with good development of the musculature of the extremities, represented by relaxed arm and calf circumference. Data reported by Arenas et al. (18) for the general Venezuelan population of similar age and by Leone et al. (19) in female swimmers, suggest the important role played by this last variable in the characterization of groups. On the other hand, as growth proceeds in

these children and youths, their upper body fat increases.

Characterizing variables were successfully identified that could separate the three maturity status groups. The highest prediction capabilities in all categories were offered by stretch stature, triceps and subscapular skinfolds, which were found to be the dominant variables, producing an impressive hit rate (75%) in prepubescent. In order to reach the highest hit rate (80%) in the aforementioned group, calf circumference, relaxed arm and body mass were also required. This set of variables were also essential to attain 83% correct classification in initial pubescent. In contrast, just stretch stature and triceps skinfold result in a hit rate of 89% for advanced pubescent. The eight variables selected by the discriminant analysis process were sufficient to attain 80%, 87% and 93% apparently correct classification of prepubescent, initial pubescent and advanced pubescent respectively.

Since discriminant analysis was taken as the decision rule, if the variables used for this analysis are available, then a simple program can be used to estimate the probability that a subject will fall into a given group, prepubescence, initial pubescence, or advanced pubescence. In practice, however, this information will very probably not be available, and so means and their confidence intervals can be used as a reference for estimating maturation categories.

While coaches should consider the potential genetic predisposition of young athletes in talent identification and development programs, it is also necessary that maturity indicators, which are partly related to the endocrinological changes of puberty, be taken into account. Compelling data (20), (21), (22), (23) indicate that biological age rather than chronological age is a better predictor of physical performance, since a given bone age can correspond to a wide range of chronological ages.

CONCLUSIONS

There can be no doubt that the growth and maturation process, together with other sets of variables, is essential for the identification of athletic talents as changes occur, both in physical structure and performance. It may be concluded from this study that - as determined in this particular sample - certain anthropometrical variables have a specific weight to separate groups of maturity status. The present paper provide criteria that can guide coaches in the classification of swimmers by certain biological characteristics, which will bring the training they provide more into line with the biological status of specific athletes.

REFERENCES

- Bale P. The functional performance of children in relation to growth, maturation and exercise. *Sports Med* 1992; 13 (3): 151-9.
- Bass Shona L. The prepubertal years. *Sports Med* 2000; 30 (2): 74-8,
- Beunen G, Malina R. Growth and biological maturation: relevance to athletic performance. In: O Bar-Or, editor. *The child and adolescent athlete*. Oxford: Blackwell Science; 1996 p. 3-24
- Baxter-Jones, A.D. Growth and development of young athletes. *Sports Med* 1995; 20: 59-64.
- Landers G.J, B.A. Blanksby, T.R. Ackland and D.A. Smith. Kinanthropometric differences between world championship senior and junior elite triathletes. 1999. Department of Human Movement and Exercise Science, The University of Western Australia, Perth, Australia. Available on: <<http://www.ausport.gov.au/fulltext/triathlon/landers.blanksby.ackland.smith.pdf>> 2004 04/06/ 1:30pm
- Pérez B, Landaeta-Jiménez M, editors. *Perfil biológico y nutricional de los nadadores del estado Miranda*. Caracas: Ediciones del Vicerrectorado Académico UCV; 2004.
- International Standards for Anthropometric Assessment. (ISAK) Australia: International Society for the Advancement of Kinanthropometry ; 2001.
- Norton K., T. Olds .*Anthropometrica: A textbook of body measurements for sports and health courses*. Sydney Australia. University of New South Wales Press; 1996.
- Gore Ch, Olds T, Carter L, Norton K. *Accreditation in anthropometry. A curriculum guide*. National Sport Research Center. Australian Sport Commission; 1998.
- Pérez B, Ortega de Mancera A, Ledezma T. and Salazar L. M. *Metodología General*. In: Pérez B.M, M. Landaeta-Jiménez editors. *Perfil biológico y estado nutricional en nadadores del estado Miranda*. Caracas: Universidad Central de Venezuela. 2004. p.19-51.
- Tanner J.M. *Growth at adolescence*. 2nd ed. Oxford. Blackwell Scientific Publications; 1962.
- Nicoletti I. Condizionamenti della stature e standard condizionati. In: *La Crescita del Bambino Italiano*. Centro Studi Auxologici. Firenze, Italia:1992. p.23-40
- Marshall W.A., Tanner J.M. Variation in the pattern of pubertal changes in boys. *Arch Dis Child* 1970; 45, 13-23.
- Huberty C. *Applied discriminant analysis*. New York: Wiley; 1994.
- Malina RM, Bielicki T. Retrospective longitudinal growth study of boys and girls active in sport. *Acta Paediatr* 1996; 85: 570-6.
- Macías-Tomei C, Izaguirre-Espinoza I, López-Blanco M. Maduración sexual y ósea según ritmo en niños y jóvenes del Estudio Longitudinal de Caracas. *An Ven Nutr* 2000; 13(1): 188-195
- Peña Reyes, M. E. and Malina, R. M. Growth and maturity profile of youth swimmers in Mexico. In: Manuel Coelho E Silva, Robert M. Malina editors. *Children and youth in organized sports*. Coimbra: Coimbra University Press; 2004. p 222-30.
- Arenas O, Pérez B.M., and Castillo T. Variables antropométricas y su potencial discriminatorio en grupos de individuos estratificados por sexo y edad. *Acta Científica Venezolana* 1988; 39: 375-9.
- Leone, M., Lariviere, G. and Comtois A Discriminant analysis of anthropometric and biomotor variables among elite adolescent female athletes in four sports. *J Sports Sci*. 2002;20(7):i6, p.443
- Malina, R. Physical growth and biological maturation of young athletes. *Exerc Sport Sci Rev* 1994;22:389-433.
- Geithner, C.A., Woynarowska B, Malina R. The adolescent spurt and sexual maturity in girls active and not active in sport. *Ann Hum Biol* 1998; 25: 413-23.
- Jones MA, Hitchen PJ, Stratton G. The importance of considering biological maturity when assessing physical fitness measures in girls and boys aged 10 to 16 years. *Ann Hum Biol* 2000; 27 (1):57-65
- Malina R. Growth and maturity status of children in organized sports. In: Nicoletti I, Benso L, Gilli G, editors. *Physiological and Pathological Auxology*. Firenze Italy: Edizioni Centro Studi Auxologici; 2004. p.595-605.

ACKNOWLEDGMENTS

This research was supported by the Consejo de Desarrollo Científico y Humanístico of the Central University of Venezuela (PG-05-30-474100). The authors would like to thank to anthropologist Marínés Salazar L, research assistant of the Department of Bioanthropology, Physical Activity and Health, who was involved in different phases of the project from which this data was derived. Additionally, the authors are grateful to Dr. J.E.L. Carter for criticism and suggestions to enhance the quality of this paper.

Endereço para correspondência:

Betty M. Pérez
El Marqués. Unidad Residencial El Portón.
Calles "A" y "B". Quinta Mariusa. Caracas, 1070. Venezuela.
E-mail: mariusa@telcel.net.ve

Recebido em 23/01/06

Revisado em 03/02/06

Aprovado em 10/02/06